

Generalized (anti) Yetter-Drinfeld modules as components of a braided T-category

Florin Panaite*

Institute of Mathematics of the Romanian Academy
PO-Box 1-764, RO-014700 Bucharest, Romania
e-mail: Florin.Panaite@imar.ro

Mihai D. Staic†

SUNY at Buffalo
Amherst, NY 14260-2900, USA
e-mail: mdstaic@buffalo.edu

Abstract

If H is a Hopf algebra with bijective antipode and $\alpha, \beta \in \text{Aut}_{\text{Hopf}}(H)$, we introduce a category ${}_H\mathcal{YD}^H(\alpha, \beta)$, generalizing both Yetter-Drinfeld modules and anti-Yetter-Drinfeld modules. We construct a braided T-category $\mathcal{YD}(H)$ having all the categories ${}_H\mathcal{YD}^H(\alpha, \beta)$ as components, which if H is finite dimensional coincides with the representations of a certain quasitriangular T-coalgebra $DT(H)$ that we construct. We also prove that if (α, β) admits a so-called pair in involution, then ${}_H\mathcal{YD}^H(\alpha, \beta)$ is isomorphic to the category of usual Yetter-Drinfeld modules ${}_H\mathcal{YD}^H$.

Introduction

Let H be a Hopf algebra with bijective antipode S and $\alpha, \beta \in \text{Aut}_{\text{Hopf}}(H)$. We introduce the concept of an (α, β) -Yetter-Drinfeld module, as being a left H -module right H -comodule M with the following compatibility condition:

$$(h \cdot m)_{(0)} \otimes (h \cdot m)_{(1)} = h_2 \cdot m_{(0)} \otimes \beta(h_3)m_{(1)}\alpha(S^{-1}(h_1)).$$

This concept is a generalization of three kinds of objects appeared in the literature. Namely, for $\alpha = \beta = id_H$, we obtain the usual Yetter-Drinfeld modules; for $\alpha = S^2$, $\beta = id_H$, we obtain the so-called anti-Yetter-Drinfeld modules, introduced in [7], [8], [10] as coefficients for the cyclic cohomology of Hopf algebras defined by Connes and Moscovici in [5], [6]; finally, an (id_H, β) -Yetter-Drinfeld module is a generalization of the object H_β defined in [4], which has the property that, if H is finite dimensional, then the map $\beta \mapsto \text{End}(H_\beta)$ gives a group anti-homomorphism from $\text{Aut}_{\text{Hopf}}(H)$ to the Brauer group of H .

It is natural to expect that (α, β) -Yetter-Drinfeld modules have some properties resembling the

*Research partially supported by the programme CERES of the Romanian Ministry of Education and Research, contract no. 4-147/2004.

†Permanent address: Institute of Mathematics of the Romanian Academy, PO-Box 1-764, RO-014700 Bucharest, Romania.

ones of the three kinds of objects we mentioned. We will see some of these properties in this paper (others will be given in a subsequent one), namely the ones directed to our main aim here, which is the following: if we denote by ${}_H\mathcal{YD}^H(\alpha, \beta)$ the category of (α, β) -Yetter-Drinfeld modules and we define $\mathcal{YD}(H)$ as the disjoint union of all these categories, then we can organize $\mathcal{YD}(H)$ as a braided T-category (or braided crossed group-category, in the original terminology of Turaev, see [16]) over the group $G = \text{Aut}_{\text{Hopf}}(H) \times \text{Aut}_{\text{Hopf}}(H)$ with multiplication $(\alpha, \beta) * (\gamma, \delta) = (\alpha\gamma, \delta\gamma^{-1}\beta\gamma)$. We also prove that the subcategory $\mathcal{YD}(H)_{fd}$ consisting of finite dimensional objects has left and right dualities, and that, if H is finite dimensional, then $\mathcal{YD}(H)$ coincides with the representations of a certain quasitriangular T-coalgebra $DT(H)$ that we construct. Our second aim is to prove that, if $\alpha, \beta \in \text{Aut}_{\text{Hopf}}(H)$ such that there exists a so-called *pair in involution* (f, g) corresponding to (α, β) , then ${}_H\mathcal{YD}^H(\alpha, \beta)$ is isomorphic to ${}_H\mathcal{YD}^H$. This result is independent on the theory concerning $\mathcal{YD}(H)$, but we can give it a very short proof using the results obtained during the construction of $\mathcal{YD}(H)$.

1 Preliminaries

We work over a ground field k . All algebras, linear spaces, etc. will be over k ; unadorned $\otimes$ means $\otimes_k$. Unless otherwise stated, H will denote a Hopf algebra with bijective antipode S . We will use the versions of Sweedler's sigma notation: $\Delta(h) = h_1 \otimes h_2$ or $\Delta(h) = h_{(1)} \otimes h_{(2)}$. For unexplained concepts and notation about Hopf algebras we refer to [11], [12], [13], [15]. By $\alpha, \beta, \gamma, \dots$ we will usually denote Hopf automorphisms of H .

Let A be an H -bicomodule algebra, with comodule structures $A \rightarrow A \otimes H$, $a \mapsto a_{<0>} \otimes a_{<1>}$ and $A \rightarrow H \otimes A$, $a \mapsto a_{[-1]} \otimes a_{[0]}$, and denote, for $a \in A$,

$$a_{\{-1\}} \otimes a_{\{0\}} \otimes a_{\{1\}} = a_{<0>[-1]} \otimes a_{<0>[0]} \otimes a_{<1>} = a_{[-1]} \otimes a_{[0]<0>} \otimes a_{[0]<1>}$$

as an element in $H \otimes A \otimes H$. We can consider the *Yetter-Drinfeld datum* (H, A, H) as in [3] (the second H is regarded as an H -bimodule coalgebra), and the Yetter-Drinfeld category ${}_A\mathcal{YD}(H)^H$, whose objects are k -modules M endowed with a left A -action (denoted by $a \otimes m \mapsto a \cdot m$) and a right H -coaction (denoted by $m \mapsto m_{(0)} \otimes m_{(1)}$) satisfying the equivalent compatibility conditions

$$(a \cdot m)_{(0)} \otimes (a \cdot m)_{(1)} = a_{\{0\}} \cdot m_{(0)} \otimes a_{\{1\}} m_{(1)} S^{-1}(a_{\{-1\}}), \quad (1.1)$$

$$a_{<0>} \cdot m_{(0)} \otimes a_{<1>} m_{(1)} = (a_{[0]} \cdot m)_{(0)} \otimes (a_{[0]} \cdot m)_{(1)} a_{[-1]}, \quad (1.2)$$

for all $a \in A$ and $m \in M$.

Recall now from [9] the construction of the (left) *diagonal crossed product* $H^* \bowtie A$, which is an associative algebra constructed on $H^* \otimes A$, with multiplication given by

$$(p \bowtie a)(q \bowtie b) = p(a_{\{-1\}} \rightharpoonup q \leftharpoonup S^{-1}(a_{\{1\}})) \bowtie a_{\{0\}} b, \quad (1.3)$$

for all $a, b \in A$ and $p, q \in H^*$, and with unit $\varepsilon_H \bowtie 1_A$. Here $\rightharpoonup$ and $\leftharpoonup$ are the regular actions of H on H^* given by $(h \rightharpoonup p)(l) = p(lh)$ and $(p \leftharpoonup h)(l) = p(hl)$ for all $h, l \in H$ and $p \in H^*$.

If H is finite dimensional, we can consider the Drinfeld double $D(H)$, which is a quasitriangular Hopf algebra realized on $H^* \otimes H$; its coalgebra structure is $H^{*cop} \otimes H$ and the algebra structure is just $H^* \bowtie H$, that is

$$(p \bowtie h)(q \bowtie l) = p(h_1 \rightharpoonup q \leftharpoonup S^{-1}(h_3)) \bowtie h_2 l, \quad (1.4)$$

for all $p, q \in H^*$ and $h, l \in H$.

The diagonal crossed product $H^* \bowtie A$ becomes a $D(H)$ -bicomodule algebra, with structures

$$\begin{aligned} H^* \bowtie A &\rightarrow (H^* \bowtie A) \otimes D(H), \quad p \bowtie a \mapsto (p_2 \bowtie a_{<0>}) \otimes (p_1 \otimes a_{<1>}), \\ H^* \bowtie A &\rightarrow D(H) \otimes (H^* \bowtie A), \quad p \bowtie a \mapsto (p_2 \otimes a_{[-1]}) \otimes (p_1 \bowtie a_{[0]}), \end{aligned}$$

for all $p \in H^*$ and $a \in A$, see [9].

In the case when H is finite dimensional, by results in [1], [3] it follows that the category ${}_A\mathcal{YD}(H)^H$ is isomorphic to the category ${}_{H^* \bowtie A}\mathcal{M}$ of left modules over $H^* \bowtie A$.

2 (α, β) -Yetter-Drinfeld modules

Definition 2.1 Let $\alpha, \beta \in \text{Aut}_{\text{Hopf}}(H)$. An (α, β) -Yetter-Drinfeld module over H is a vector space M , such that M is a left H -module (with notation $h \cdot m \mapsto h \cdot m$) and a right H -comodule (with notation $M \rightarrow M \otimes H$, $m \mapsto m_{(0)} \otimes m_{(1)}$) with the following compatibility condition:

$$(h \cdot m)_{(0)} \otimes (h \cdot m)_{(1)} = h_2 \cdot m_{(0)} \otimes \beta(h_3)m_{(1)}\alpha(S^{-1}(h_1)), \quad (2.1)$$

for all $h \in H$ and $m \in M$. We denote by ${}_H\mathcal{YD}^H(\alpha, \beta)$ the category of (α, β) -Yetter-Drinfeld modules, morphisms being the H -linear H -colinear maps.

Remark 2.2 As for usual Yetter-Drinfeld modules, one can see that (2.1) is equivalent to

$$h_1 \cdot m_{(0)} \otimes \beta(h_2)m_{(1)} = (h_2 \cdot m)_{(0)} \otimes (h_2 \cdot m)_{(1)}\alpha(h_1). \quad (2.2)$$

Example 2.3 For $\alpha = \beta = \text{id}_H$, we have ${}_H\mathcal{YD}^H(\text{id}, \text{id}) = {}_H\mathcal{YD}^H$, the usual category of (left-right) Yetter-Drinfeld modules.

Example 2.4 For $\alpha = S^2$, $\beta = \text{id}_H$, the compatibility condition (2.1) becomes

$$(h \cdot m)_{(0)} \otimes (h \cdot m)_{(1)} = h_2 \cdot m_{(0)} \otimes h_3m_{(1)}S(h_1), \quad (2.3)$$

hence ${}_H\mathcal{YD}^H(S^2, \text{id})$ is the category of anti-Yetter-Drinfeld modules defined in [7], [8], [10].

Example 2.5 For $\beta \in \text{Aut}_{\text{Hopf}}(H)$, define H_β as in [4], that is $H_\beta = H$, with regular right H -comodule structure and left H -module structure given by $h \cdot h' = \beta(h_2)h'S^{-1}(h_1)$, for all $h, h' \in H$. It was noticed in [4] that H_β satisfies a certain compatibility condition, which actually says that $H_\beta \in {}_H\mathcal{YD}^H(\text{id}, \beta)$. More generally, if $\alpha, \beta \in \text{Aut}_{\text{Hopf}}(H)$, define $H_{\alpha, \beta}$ as follows: $H_{\alpha, \beta} = H$, with regular right H -comodule structure and left H -module structure given by $h \cdot h' = \beta(h_2)h'\alpha(S^{-1}(h_1))$, for $h, h' \in H$. Then one can check that $H_{\alpha, \beta} \in {}_H\mathcal{YD}^H(\alpha, \beta)$.

Example 2.6 Take l an integer and define $\alpha_l = S^{2l} \in \text{Aut}_{\text{Hopf}}(H)$. The compatibility in ${}_H\mathcal{YD}^H(S^{2l}, \text{id})$ becomes

$$(h \cdot m)_{(0)} \otimes (h \cdot m)_{(1)} = h_2 \cdot m_{(0)} \otimes h_3m_{(1)}S^{2l-1}(h_1). \quad (2.4)$$

An object in ${}_H\mathcal{YD}^H(S^{2l}, \text{id})$ will be called an l - $\mathcal{YD}$ -module. Hence, a 0- $\mathcal{YD}$ -module is a Yetter-Drinfeld module and a 1- $\mathcal{YD}$ -module is an anti-Yetter-Drinfeld module. The right-left version of l - $\mathcal{YD}$ -modules has been introduced in [14].

Example 2.7 Let $\alpha, \beta \in \text{Aut}_{H\text{opf}}(H)$ and assume that there exist an algebra map $f : H \rightarrow k$ and a group-like element $g \in H$ such that

$$\alpha(h) = g^{-1}f(h_1)\beta(h_2)f(S(h_3))g, \quad \forall h \in H. \quad (2.5)$$

Then one can check that $k \in {}_H\mathcal{YD}^H(\alpha, \beta)$, with structures $h \cdot 1 = f(h)$ and $1 \mapsto 1 \otimes g$. More generally, if V is any vector space, then $V \in {}_H\mathcal{YD}^H(\alpha, \beta)$, with structures $h \cdot v = f(h)v$ and $v \mapsto v \otimes g$, for all $h \in H$ and $v \in V$.

Definition 2.8 If $\alpha, \beta \in \text{Aut}_{H\text{opf}}(H)$ such that there exist f, g as in Example 2.7, we will say that (f, g) is a pair in involution corresponding to (α, β) (in analogy with the concept of modular pair in involution due to Connes and Moscovici) and the (α, β) -Yetter-Drinfeld modules k and V constructed in Example 2.7 will be denoted by ${}_fk^g$ and respectively ${}_fV^g$.

As an example, if $\alpha \in \text{Aut}_{H\text{opf}}(H)$, then $(\varepsilon, 1)$ is a pair in involution corresponding to (α, α) .

Let $\alpha, \beta \in \text{Aut}_{H\text{opf}}(H)$. We define an H -bicomodule algebra $H(\alpha, \beta)$ as follows: $H(\alpha, \beta) = H$ as algebra, with comodule structures

$$\begin{aligned} H(\alpha, \beta) &\rightarrow H \otimes H(\alpha, \beta), & h &\mapsto h_{[-1]} \otimes h_{[0]} = \alpha(h_1) \otimes h_2, \\ H(\alpha, \beta) &\rightarrow H(\alpha, \beta) \otimes H, & h &\mapsto h_{<0>} \otimes h_{<1>} = h_1 \otimes \beta(h_2). \end{aligned}$$

Then we can consider the Yetter-Drinfeld datum $(H, H(\alpha, \beta), H)$ and the Yetter-Drinfeld modules over it, ${}_{H(\alpha, \beta)}\mathcal{YD}(H)^H$.

Proposition 2.9 ${}_H\mathcal{YD}^H(\alpha, \beta) = {}_{H(\alpha, \beta)}\mathcal{YD}(H)^H$.

Proof. It is easy to see that the compatibility conditions for the two categories are the same. $\square$

In particular, the category of anti-Yetter-Drinfeld modules coincides with ${}_{H(S^2, id)}\mathcal{YD}(H)^H$, which improves the remark in [7] that anti-Yetter-Drinfeld modules are entwined modules.

Consider now the diagonal crossed product $A(\alpha, \beta) = H^* \bowtie H(\alpha, \beta)$, whose multiplication is

$$(p \bowtie h)(q \bowtie l) = p(\alpha(h_1) \rightharpoonup q \leftarrow S^{-1}(\beta(h_3))) \bowtie h_2l, \quad (2.6)$$

for all $p, q \in H^*$ and $h, l \in H$. For $\alpha = \beta = id$ we get $A(id, id) = D(H)$; for $\alpha = S^2$ and $\beta = id$, the multiplication in $A(S^2, id)$ is

$$(p \bowtie h)(q \bowtie l) = p(S^2(h_1) \rightharpoonup q \leftarrow S^{-1}(h_3)) \bowtie h_2l, \quad (2.7)$$

hence $A(S^2, id)$ coincides with the algebra $A(H)$ defined in [7].

Assume now that H is finite dimensional; then $A(\alpha, \beta)$ becomes a $D(H)$ -bicomodule algebra, with structures

$$\begin{aligned} H^* \bowtie H(\alpha, \beta) &\rightarrow (H^* \bowtie H(\alpha, \beta)) \otimes D(H), & p \bowtie h &\mapsto (p_2 \bowtie h_1) \otimes (p_1 \otimes \beta(h_2)), \\ H^* \bowtie H(\alpha, \beta) &\rightarrow D(H) \otimes (H^* \bowtie H(\alpha, \beta)), & p \bowtie h &\mapsto (p_2 \otimes \alpha(h_1)) \otimes (p_1 \bowtie h_2). \end{aligned}$$

In particular, $A(H)$ becomes a $D(H)$ -bicomodule algebra, improving the remark in [7] that $A(H)$ is a right $D(H)$ -comodule algebra. Since H is finite dimensional, we have an isomorphism of categories ${}_{H(\alpha, \beta)}\mathcal{YD}(H)^H \simeq {}_{H^* \bowtie H(\alpha, \beta)}\mathcal{M}$, hence ${}_H\mathcal{YD}^H(\alpha, \beta) \simeq {}_{H^* \bowtie H(\alpha, \beta)}\mathcal{M}$ (for $\alpha = S^2$,

$\beta = id$ we recover the result in [7] that the category of anti-Yetter-Drinfeld modules is isomorphic to ${}_{A(H)}\mathcal{M}$. The correspondence is given as follows. If $M \in {}_H\mathcal{YD}^H(\alpha, \beta)$, then $M \in {}_{H^*\bowtie H(\alpha, \beta)}\mathcal{M}$ with structure

$$(p \bowtie h) \cdot m = p((h \cdot m)_{(1)})(h \cdot m)_{(0)}.$$

Conversely, if $M \in {}_{H^*\bowtie H(\alpha, \beta)}\mathcal{M}$, then $M \in {}_H\mathcal{YD}^H(\alpha, \beta)$ with structures

$$\begin{aligned} h \cdot m &= (\varepsilon \bowtie h) \cdot m, \\ m &\mapsto m_{(0)} \otimes m_{(1)} = (e^i \bowtie 1) \cdot m \otimes e_i, \end{aligned}$$

where $\{e_i\}, \{e^i\}$ are dual bases in H and H^* .

3 A braided T-category $\mathcal{YD}(H)$

Let $\alpha, \beta \in \text{Aut}_{\text{Hopf}}(H)$ and consider the objects H_α, H_β as in Example 2.5. In [4] was considered the object $M = H_\alpha \otimes H_\beta$, with the following structures:

$$\begin{aligned} h \cdot (x \otimes y) &= h_1 \cdot x \otimes \alpha(h_2) \cdot y, \\ x \otimes y &\mapsto (x_1 \otimes y_1) \otimes y_2 x_2, \end{aligned}$$

for all $h, x, y \in H$, where by $\cdot$ we denoted both the actions of H on H_α and H_β given as in Example 2.5. Then it was noticed in [4] that M satisfies a compatibility condition which says that $M \in {}_H\mathcal{YD}^H(id, \beta\alpha)$.

On the other hand, it was noticed in [7] that the tensor product between an anti-Yetter-Drinfeld module and a Yetter-Drinfeld module becomes an anti-Yetter-Drinfeld module.

The next result can be seen as a generalization of both these facts.

Proposition 3.1 *If $M \in {}_H\mathcal{YD}^H(\alpha, \beta)$, $N \in {}_H\mathcal{YD}^H(\gamma, \delta)$, then $M \otimes N \in {}_H\mathcal{YD}^H(\alpha\gamma, \delta\gamma^{-1}\beta\gamma)$, with structures:*

$$\begin{aligned} h \cdot (m \otimes n) &= \gamma(h_1) \cdot m \otimes \gamma^{-1}\beta\gamma(h_2) \cdot n, \\ m \otimes n &\mapsto (m \otimes n)_{(0)} \otimes (m \otimes n)_{(1)} = (m_{(0)} \otimes n_{(0)}) \otimes n_{(1)}m_{(1)}. \end{aligned}$$

Proof. Obviously $M \otimes N$ is a left H -module and a right H -comodule. We check now the compatibility condition. We compute:

$$\begin{aligned} &(h \cdot (m \otimes n))_{(0)} \otimes (h \cdot (m \otimes n))_{(1)} \\ &= (\gamma(h_1) \cdot m \otimes \gamma^{-1}\beta\gamma(h_2) \cdot n)_{(0)} \otimes (\gamma(h_1) \cdot m \otimes \gamma^{-1}\beta\gamma(h_2) \cdot n)_{(1)} \\ &= ((\gamma(h_1) \cdot m)_{(0)} \otimes (\gamma^{-1}\beta\gamma(h_2) \cdot n)_{(0)}) \otimes (\gamma^{-1}\beta\gamma(h_2) \cdot n)_{(1)}(\gamma(h_1) \cdot m)_{(1)} \\ &= (\gamma(h_1)_2 \cdot m_{(0)} \otimes \gamma^{-1}\beta\gamma(h_2)_2 \cdot n_{(0)}) \otimes \\ &\quad \otimes \delta(\gamma^{-1}\beta\gamma(h_2)_3 n_{(1)} \gamma(S^{-1}(\gamma^{-1}\beta\gamma(h_2)_1)) \beta(\gamma(h_1)_3) m_{(1)} \alpha(S^{-1}(\gamma(h_1)_1))) \\ &= (\gamma(h_2) \cdot m_{(0)} \otimes \gamma^{-1}\beta\gamma(h_5) \cdot n_{(0)}) \otimes \delta\gamma^{-1}\beta\gamma(h_6) n_{(1)} \beta\gamma(S^{-1}(h_4)) \beta\gamma(h_3) m_{(1)} \alpha(S^{-1}(\gamma(h_1))) \\ &= (\gamma(h_2) \cdot m_{(0)} \otimes \gamma^{-1}\beta\gamma(h_3) \cdot n_{(0)}) \otimes \delta\gamma^{-1}\beta\gamma(h_4) n_{(1)} m_{(1)} \alpha\gamma(S^{-1}(h_1)) \\ &= h_2 \cdot (m_{(0)} \otimes n_{(0)}) \otimes \delta\gamma^{-1}\beta\gamma(h_3) n_{(1)} m_{(1)} \alpha\gamma(S^{-1}(h_1)) \\ &= h_2 \cdot (m \otimes n)_{(0)} \otimes \delta\gamma^{-1}\beta\gamma(h_3)(m \otimes n)_{(1)} \alpha\gamma(S^{-1}(h_1)), \end{aligned}$$

that is $M \otimes N \in {}_H\mathcal{YD}^H(\alpha\gamma, \delta\gamma^{-1}\beta\gamma)$. $\square$

Note that, if $M \in {}_H\mathcal{YD}^H(\alpha, \beta)$, $N \in {}_H\mathcal{YD}^H(\gamma, \delta)$ and $P \in {}_H\mathcal{YD}^H(\mu, \nu)$, then $(M \otimes N) \otimes P = M \otimes (N \otimes P)$ as objects in ${}_H\mathcal{YD}^H(\alpha\gamma\mu, \nu\mu^{-1}\delta\gamma^{-1}\beta\gamma\mu)$.

Denote $G = \text{Aut}_{\text{Hopf}}(H) \times \text{Aut}_{\text{Hopf}}(H)$, a group with multiplication

$$(\alpha, \beta) * (\gamma, \delta) = (\alpha\gamma, \delta\gamma^{-1}\beta\gamma) \quad (3.1)$$

(the unit is (id, id) and $(\alpha, \beta)^{-1} = (\alpha^{-1}, \alpha\beta^{-1}\alpha^{-1})$).

Proposition 3.2 *Let $N \in {}_H\mathcal{YD}^H(\gamma, \delta)$ and $(\alpha, \beta) \in G$. Define ${}^{(\alpha, \beta)}N = N$ as vector space, with structures*

$$\begin{aligned} h \rightharpoonup n &= \gamma^{-1}\beta\gamma\alpha^{-1}(h) \cdot n, \\ n &\mapsto n_{<0>} \otimes n_{<1>} = n_{(0)} \otimes \alpha\beta^{-1}(n_{(1)}). \end{aligned}$$

Then ${}^{(\alpha, \beta)}N \in {}_H\mathcal{YD}^H(\alpha\gamma\alpha^{-1}, \alpha\beta^{-1}\delta\gamma^{-1}\beta\gamma\alpha^{-1}) = {}_H\mathcal{YD}^H((\alpha, \beta) * (\gamma, \delta) * (\alpha, \beta)^{-1})$.

Proof. Obviously ${}^{(\alpha, \beta)}N$ is a left H -module and right H -comodule, so we check the compatibility condition. We compute:

$$\begin{aligned} (h \rightharpoonup n)_{<0>} \otimes (h \rightharpoonup n)_{<1>} &= (\gamma^{-1}\beta\gamma\alpha^{-1}(h) \cdot n)_{(0)} \otimes \alpha\beta^{-1}((\gamma^{-1}\beta\gamma\alpha^{-1}(h) \cdot n)_{(1)}) \\ &= \gamma^{-1}\beta\gamma\alpha^{-1}(h_2) \cdot n_{(0)} \otimes \alpha\beta^{-1}(\delta\gamma^{-1}\beta\gamma\alpha^{-1}(h_3)n_{(1)}\gamma\gamma^{-1}\beta\gamma\alpha^{-1}(S^{-1}(h_1))) \\ &= \gamma^{-1}\beta\gamma\alpha^{-1}(h_2) \cdot n_{(0)} \otimes \alpha\beta^{-1}\delta\gamma^{-1}\beta\gamma\alpha^{-1}(h_3)\alpha\beta^{-1}(n_{(1)})\alpha\gamma\alpha^{-1}(S^{-1}(h_1)) \\ &= h_2 \rightharpoonup n_{(0)} \otimes \alpha\beta^{-1}\delta\gamma^{-1}\beta\gamma\alpha^{-1}(h_3)n_{<1>}\alpha\gamma\alpha^{-1}(S^{-1}(h_1)), \end{aligned}$$

that is ${}^{(\alpha, \beta)}N \in {}_H\mathcal{YD}^H(\alpha\gamma\alpha^{-1}, \alpha\beta^{-1}\delta\gamma^{-1}\beta\gamma\alpha^{-1})$. $\square$

Remark 3.3 *Let $M \in {}_H\mathcal{YD}^H(\alpha, \beta)$, $N \in {}_H\mathcal{YD}^H(\gamma, \delta)$ and $(\mu, \nu) \in G$. Then we have*

$${}^{(\alpha, \beta)*(\mu, \nu)}N = {}^{(\alpha, \beta)}({}^{(\mu, \nu)}N)$$

as objects in ${}_H\mathcal{YD}^H(\alpha\mu\gamma\mu^{-1}\alpha^{-1}, \alpha\beta^{-1}\mu\nu^{-1}\delta\gamma^{-1}\nu\mu^{-1}\beta\mu\gamma\mu^{-1}\alpha^{-1})$, and

$${}^{(\mu, \nu)}(M \otimes N) = {}^{(\mu, \nu)}M \otimes {}^{(\mu, \nu)}N$$

as objects in ${}_H\mathcal{YD}^H(\mu\alpha\gamma\mu^{-1}, \mu\nu^{-1}\delta\gamma^{-1}\beta\alpha^{-1}\nu\alpha\gamma\mu^{-1})$.

Proposition 3.4 *Let $M \in {}_H\mathcal{YD}^H(\alpha, \beta)$ and $N \in {}_H\mathcal{YD}^H(\gamma, \delta)$. Define ${}^M N = {}^{(\alpha, \beta)}N$ as object in ${}_H\mathcal{YD}^H((\alpha, \beta) * (\gamma, \delta) * (\alpha, \beta)^{-1})$. Define the map*

$$c_{M, N} : M \otimes N \rightarrow {}^M N \otimes M, \quad c_{M, N}(m \otimes n) = n_{(0)} \otimes \beta^{-1}(n_{(1)}) \cdot m.$$

Then $c_{M, N}$ is H -linear H -colinear and satisfies the conditions (for $P \in {}_H\mathcal{YD}^H(\mu, \nu)$):

$$c_{M \otimes N, P} = (c_{M, N} \otimes id_P) \circ (id_M \otimes c_{N, P}), \quad (3.2)$$

$$c_{M, N \otimes P} = (id_M \otimes c_{N, P}) \circ (c_{M, N} \otimes id_P). \quad (3.3)$$

Moreover, if $M \in {}_H\mathcal{YD}^H(\alpha, \beta)$, $N \in {}_H\mathcal{YD}^H(\gamma, \delta)$ and $(\mu, \nu) \in G$, then $c_{({}^{(\mu, \nu)}M), ({}^{(\mu, \nu)}N)} = c_{M, N}$.

Proof. We prove that $c_{M,N}$ is H -linear. We compute:

$$\begin{aligned}
c_{M,N}(h \cdot (m \otimes n)) &= c_{M,N}(\gamma(h_1) \cdot m \otimes \gamma^{-1}\beta\gamma(h_2) \cdot n) \\
&= (\gamma^{-1}\beta\gamma(h_2) \cdot n)_{(0)} \otimes \beta^{-1}((\gamma^{-1}\beta\gamma(h_2) \cdot n)_{(1)})\gamma(h_1) \cdot m \\
&= \gamma^{-1}\beta\gamma(h_2)_2 \cdot n_{(0)} \otimes \beta^{-1}(\delta(\gamma^{-1}\beta\gamma(h_2)_3)n_{(1)}\gamma(S^{-1}((\gamma^{-1}\beta\gamma(h_2))_1)))\gamma(h_1) \cdot m \\
&= \gamma^{-1}\beta\gamma(h_3) \cdot n_{(0)} \otimes \beta^{-1}\delta\gamma^{-1}\beta\gamma(h_4)\beta^{-1}(n_{(1)})\gamma(S^{-1}(h_2))\gamma(h_1) \cdot m \\
&= \gamma^{-1}\beta\gamma(h_1) \cdot n_{(0)} \otimes \beta^{-1}\delta\gamma^{-1}\beta\gamma(h_2)\beta^{-1}(n_{(1)}) \cdot m,
\end{aligned}$$

$$\begin{aligned}
h \cdot c_{M,N}(m \otimes n) &= h \cdot (n_{(0)} \otimes \beta^{-1}(n_{(1)}) \cdot m) \\
&= \alpha(h_1) \rightharpoonup n_{(0)} \otimes \alpha^{-1}\alpha\beta^{-1}\delta\gamma^{-1}\beta\gamma\alpha^{-1}\alpha(h_2)\beta^{-1}(n_{(1)}) \cdot m \\
&= \gamma^{-1}\beta\gamma\alpha^{-1}\alpha(h_1) \cdot n_{(0)} \otimes \beta^{-1}\delta\gamma^{-1}\beta\gamma(h_2)\beta^{-1}(n_{(1)}) \cdot m \\
&= \gamma^{-1}\beta\gamma(h_1) \cdot n_{(0)} \otimes \beta^{-1}\delta\gamma^{-1}\beta\gamma(h_2)\beta^{-1}(n_{(1)}) \cdot m,
\end{aligned}$$

so the two terms are equal. The fact that $c_{M,N}$ is H -colinear is similar and left to the reader. We prove now (3.2). First note that, due to Remark 3.3, we have ${}^M(NP) = {}^{M \otimes N}P$ and ${}^M(N \otimes P) = {}^M N \otimes {}^M P$. We compute:

$$\begin{aligned}
(c_{M, {}^N P} \otimes id_N) \circ (id_M \otimes c_{N,P})(m \otimes n \otimes p) &= c_{M, {}^N P}(m \otimes p_{(0)}) \otimes \delta^{-1}(p_{(1)}) \cdot n \\
&= p_{(0) < 0 >} \otimes \beta^{-1}(p_{(0) < 1 >}) \cdot m \otimes \delta^{-1}(p_{(1)}) \cdot n \\
&= p_{(0)(0)} \otimes \beta^{-1}\gamma\delta^{-1}(p_{(0)(1)}) \cdot m \otimes \delta^{-1}(p_{(1)}) \cdot n \\
&= p_{(0)} \otimes \beta^{-1}\gamma\delta^{-1}(p_{(1)_1}) \cdot m \otimes \delta^{-1}(p_{(1)_2}) \cdot n,
\end{aligned}$$

$$\begin{aligned}
c_{M \otimes N, P}(m \otimes n \otimes p) &= p_{(0)} \otimes \gamma^{-1}\beta^{-1}\gamma\delta^{-1}(p_{(1)}) \cdot (m \otimes n) \\
&= p_{(0)} \otimes \gamma\gamma^{-1}\beta^{-1}\gamma\delta^{-1}(p_{(1)_1}) \cdot m \otimes \gamma^{-1}\beta\gamma\gamma^{-1}\beta^{-1}\gamma\delta^{-1}(p_{(1)_2}) \cdot n \\
&= p_{(0)} \otimes \beta^{-1}\gamma\delta^{-1}(p_{(1)_1}) \cdot m \otimes \delta^{-1}(p_{(1)_2}) \cdot n,
\end{aligned}$$

and we are done. The proof of (3.3) is easier and left to the reader, and similarly the last statement of the Proposition. $\square$

Note that $c_{M,N}$ is bijective with inverse $c_{M,N}^{-1}(n \otimes m) = \beta^{-1}(S(n_{(1)})) \cdot m \otimes n_{(0)}$.

We are ready now to introduce the desired braided T-category (we use terminology as in [18]; for the subject of Turaev categories, see also the original paper of Turaev [16] and [2], [17]).

Define $\mathcal{YD}(H)$ as the disjoint union of all ${}_H\mathcal{YD}^H(\alpha, \beta)$, with $(\alpha, \beta) \in G$ (hence the component of the unit is just ${}_H\mathcal{YD}^H$). If we endow $\mathcal{YD}(H)$ with tensor product as in Proposition 3.1, then it becomes a strict monoidal category with unit k as object in ${}_H\mathcal{YD}^H$ (with trivial structures). The group homomorphism $\varphi : G \rightarrow \text{aut}(\mathcal{YD}(H))$, $(\alpha, \beta) \mapsto \varphi_{(\alpha, \beta)}$, is given on components as

$$\varphi_{(\alpha, \beta)} : {}_H\mathcal{YD}^H(\gamma, \delta) \rightarrow {}_H\mathcal{YD}^H((\alpha, \beta) * (\gamma, \delta) * (\alpha, \beta)^{-1}), \quad \varphi_{(\alpha, \beta)}(N) = {}^{(\alpha, \beta)}N,$$

and the functor $\varphi_{(\alpha, \beta)}$ acts as identity on morphisms. The braiding in $\mathcal{YD}(H)$ is given by the family $\{c_{M,N}\}$. As a consequence of the above results, we obtain:

Theorem 3.5 $\mathcal{YD}(H)$ is a braided T-category over G .

We consider now the problem of existence of left and right dualities.

Proposition 3.6 *Let $M \in {}_H\mathcal{YD}^H(\alpha, \beta)$ and assume that M is finite dimensional. Then $M^* = \text{Hom}(M, k)$ becomes an object in ${}_H\mathcal{YD}^H(\alpha^{-1}, \alpha\beta^{-1}\alpha^{-1})$, with $(h \cdot f)(m) = f((\beta^{-1}\alpha^{-1}S(h)) \cdot m)$ and $f_{(0)}(m) \otimes f_{(1)} = f(m_{(0)}) \otimes S^{-1}(m_{(1)})$. Moreover, the maps $b_M : k \rightarrow M \otimes M^*$, $b_M(1) = \sum_i e_i \otimes e^i$ (where $\{e_i\}$ and $\{e^i\}$ are dual bases in M and M^*) and $d_M : M^* \otimes M \rightarrow k$, $d_M(f \otimes m) = f(m)$, are morphisms in ${}_H\mathcal{YD}^H$ and we have $(id_M \otimes d_M)(b_M \otimes id_M) = id_M$ and $(d_M \otimes id_{M^*})(id_{M^*} \otimes b_M) = id_{M^*}$.*

Proof. We first prove that M^* is indeed an object in ${}_H\mathcal{YD}^H(\alpha^{-1}, \alpha\beta^{-1}\alpha^{-1})$. We compute:

$$\begin{aligned}
(h \cdot f)_{(0)}(m) \otimes (h \cdot f)_{(1)} &= (h \cdot f)(m_{(0)}) \otimes S^{-1}(m_{(1)}) \\
&= f((\beta^{-1}\alpha^{-1}S)(h) \cdot m_{(0)}) \otimes S^{-1}(m_{(1)}), \\
(h_{(2)} \cdot f_{(0)})(m) \otimes (\alpha\beta^{-1}\alpha^{-1})(h_{(3)})f_{(1)}(\alpha^{-1}S^{-1}(h_{(1)})) \\
&= f_{(0)}(\beta^{-1}\alpha^{-1}S(h_{(2)}) \cdot m) \otimes (\alpha\beta^{-1}\alpha^{-1})(h_{(3)})f_{(1)}(\alpha^{-1}S^{-1}(h_{(1)})) \\
&= f(((\beta^{-1}\alpha^{-1}S)(h_{(2)}) \cdot m)_{(0)}) \otimes (\alpha\beta^{-1}\alpha^{-1})(h_{(3)}) \\
&\quad S^{-1}(((\beta^{-1}\alpha^{-1}S)(h_{(2)}) \cdot m)_{(1)})(\alpha^{-1}S^{-1}(h_{(1)})) \\
&= f((\beta^{-1}\alpha^{-1}S)(h_{(3)}) \cdot m_{(0)}) \otimes (\alpha\beta^{-1}\alpha^{-1})(h_{(5)}) \\
&\quad S^{-1}((\alpha^{-1}S)(h_{(2)})m_{(1)}(\alpha\beta^{-1}\alpha^{-1})(h_{(4)}))(\alpha^{-1}S^{-1}(h_{(1)})) \\
&= f((\beta^{-1}\alpha^{-1}S)(h_{(3)}) \cdot m_{(0)}) \otimes (\alpha\beta^{-1}\alpha^{-1})(h_{(5)}S^{-1}(h_{(4)}))S^{-1}(m_{(1)})\alpha^{-1}(h_{(2)}S^{-1}(h_{(1)})) \\
&= f((\beta^{-1}\alpha^{-1}S)(h) \cdot m_{(0)}) \otimes S^{-1}(m_{(1)}),
\end{aligned}$$

which means that

$$(h \cdot f)_{(0)} \otimes (h \cdot f)_{(1)} = (h_{(2)} \cdot f_{(0)}) \otimes (\alpha\beta^{-1}\alpha^{-1})(h_{(3)})f_{(1)}(\alpha^{-1}S^{-1}(h_{(1)})), \quad q.e.d.$$

On k we have the trivial module and comodule structure, and with these $k \in {}_H\mathcal{YD}^H$. We want to prove that b_M and d_M are H -module maps. We compute:

$$\begin{aligned}
(h \cdot b_M(1))(m) &= (h \cdot (\sum_i e_i \otimes e^i))(m) \\
&= \sum_i \alpha^{-1}(h_{(1)}) \cdot e_i \otimes ((\alpha\beta\alpha^{-1})(h_{(2)}) \cdot e^i)(m) \\
&= \alpha^{-1}(h_{(1)}) \cdot e_i \otimes e^i((\beta^{-1}\alpha^{-1}S\alpha\beta\alpha^{-1})(h_{(2)}) \cdot m) \\
&= \sum_i \alpha^{-1}(h_{(1)}) \cdot e_i \otimes e^i((\alpha^{-1}S)(h_{(2)}) \cdot m) \\
&= \alpha^{-1}(h_{(1)}S(h_{(2)})) \cdot m \\
&= \varepsilon(h) \sum_i e_i \otimes e^i(m) \\
&= (\varepsilon(h)b_M(1))(m),
\end{aligned}$$

$$\begin{aligned}
d_M(h \cdot (f \otimes m)) &= d_M(\alpha(h_{(1)}) \cdot f \otimes \beta^{-1}(h_{(2)}) \cdot m) \\
&= (\alpha(h_{(1)}) \cdot f)(\beta^{-1}(h_{(2)}) \cdot m) \\
&= f((\beta^{-1}\alpha^{-1}S\alpha(h_{(1)}))\beta^{-1}(h_{(2)}) \cdot m) \\
&= f(\beta^{-1}(S(h_{(1)})h_{(2)}) \cdot m) \\
&= \varepsilon(h)d_M(f \otimes m).
\end{aligned}$$

Also they are H -comodule maps:

$$\begin{aligned}
((b_M(1))_{(0)} \otimes (b_M(1))_{(1)})(m) &= \sum_i (e_i)_{(0)} \otimes (e^i)_{(0)}(m) \otimes (e^i)_{(1)}(e_i)_{(1)} \\
&= \sum_i (e_i)_{(0)} \otimes (e^i)(m_{(0)}) \otimes S^{-1}(m_{(1)})(e_i)_{(1)} \\
&= m_{(0)} \otimes S^{-1}(m_{(1)_2})m_{(1)_1} \\
&= (b_M(1) \otimes 1)(m),
\end{aligned}$$

$$\begin{aligned}
d_M((f \otimes m)_{(0)}) \otimes (f \otimes m)_{(1)} &= f_{(0)}(m_{(0)}) \otimes m_{(1)}f_{(1)} \\
&= f(m_{(0)}) \otimes m_{(1)_2}S^{-1}(m_{(1)_1}) \\
&= d_M(f \otimes m) \otimes 1.
\end{aligned}$$

Finally, the last two identities $(id_M \otimes d_M)(b_M \otimes id_M) = id_M$ and $(d_M \otimes id_{M^*})(id_{M^*} \otimes b_M) = id_{M^*}$ are trivial. $\square$

Similarly, one can prove:

Proposition 3.7 *Let $M \in {}_H\mathcal{YD}^H(\alpha, \beta)$ and assume that M is finite dimensional. Then ${}^*M = Hom(M, k)$ becomes an object in ${}_H\mathcal{YD}^H(\alpha^{-1}, \alpha\beta^{-1}\alpha^{-1})$, with $(h \cdot f)(m) = f((\alpha^{-1}\beta^{-1}S^{-1}(h)) \cdot m)$ and $f_{(0)}(m) \otimes f_{(1)} = f(m_{(0)}) \otimes S(m_{(1)})$. Moreover, the maps $b_M : k \rightarrow {}^*M \otimes M$, $b_M(1) = \sum_i e^i \otimes e_i$ and $d_M : M \otimes {}^*M \rightarrow k$, $d_M(m \otimes f) = f(m)$, are morphisms in ${}_H\mathcal{YD}^H$ and we have $(d_M \otimes id_M)(id_M \otimes b_M) = id_M$ and $(id_{{}^*M} \otimes d_M)(b_M \otimes id_{{}^*M}) = id_{{}^*M}$.*

Consequently, if we consider $\mathcal{YD}(H)_{fd}$, the subcategory of $\mathcal{YD}(H)$ consisting of finite dimensional objects, we obtain:

Theorem 3.8 *$\mathcal{YD}(H)_{fd}$ is a braided T-category with left and right dualities over G , the left (respectively right) duals being given as in Proposition 3.6 (respectively Proposition 3.7).*

Assume now that H is finite dimensional. We will construct a quasitriangular T-coalgebra over G , denoted by $DT(H)$, with the property that the T-category $Rep(DT(H))$ of representations of $DT(H)$ is isomorphic to $\mathcal{YD}(H)$ as braided T-categories.

For $(\alpha, \beta) \in G$, the (α, β) -component $DT(H)_{(\alpha, \beta)}$ will be the diagonal crossed product algebra $H^* \bowtie H(\alpha, \beta)$. Define

$$\begin{aligned}
\Delta_{(\alpha, \beta), (\gamma, \delta)} : H^* \bowtie H((\alpha, \beta) * (\gamma, \delta)) &\rightarrow (H^* \bowtie H(\alpha, \beta)) \otimes (H^* \bowtie H(\gamma, \delta)), \\
\Delta_{(\alpha, \beta), (\gamma, \delta)}(p \bowtie h) &= (p_2 \bowtie \gamma(h_1)) \otimes (p_1 \bowtie \gamma^{-1}\beta\gamma(h_2)).
\end{aligned}$$

One can check, by direct computation, that these maps are algebra maps, satisfying the necessary coassociativity conditions.

The counit ε is just the counit of $DT(H)_{(id, id)} = D(H)$, the Drinfeld double of H .

For $(\alpha, \beta), (\gamma, \delta) \in G$, define now

$$\begin{aligned}
\varphi_{(\alpha, \beta)}^{(\gamma, \delta)} : H^* \bowtie H(\gamma, \delta) &\rightarrow H^* \bowtie H((\alpha, \beta) * (\gamma, \delta) * (\alpha, \beta)^{-1}), \\
\varphi_{(\alpha, \beta)}^{(\gamma, \delta)}(p \bowtie h) &= p \circ \beta\alpha^{-1} \bowtie \alpha\gamma^{-1}\beta^{-1}\gamma(h).
\end{aligned}$$

Then one can check by direct computation that these are algebra isomorphisms giving a *conjugation* (that is they are multiplicative and compatible with the comultiplications and the counit). The antipode is given, for $(\alpha, \beta) \in G$, by

$$\begin{aligned} S_{(\alpha, \beta)} : H^* \bowtie H(\alpha, \beta) &\rightarrow H^* \bowtie H((\alpha, \beta)^{-1}), \\ S_{(\alpha, \beta)}(p \bowtie h) &= (\varepsilon \bowtie \alpha\beta(S(h))) \cdot (S^{*-1}(p) \bowtie 1), \end{aligned}$$

where the multiplication $\cdot$ in the right hand side is made in $H^* \bowtie H((\alpha, \beta)^{-1})$. Finally, the universal R -matrix is given by

$$R_{(\alpha, \beta), (\gamma, \delta)} = \sum_i (\varepsilon \bowtie \beta^{-1}(e_i)) \otimes (e^i \bowtie 1) \in (H^* \bowtie H(\alpha, \beta)) \otimes (H^* \bowtie H(\gamma, \delta)),$$

for all $(\alpha, \beta), (\gamma, \delta) \in G$, where $\{e_i\}, \{e^i\}$ are dual bases in H and H^* . Thus, we have obtained:

Theorem 3.9 *$DT(H)$ is a quasitriangular T -coalgebra over G , with structure as above.*

Moreover, the structure of $DT(H)$ was constructed in such a way that, via the isomorphisms $H^* \bowtie H(\alpha, \beta) \mathcal{M} \simeq {}_H \mathcal{YD}^H(\alpha, \beta)$ from Section 2, we obtain:

Theorem 3.10 *$Rep(DT(H))$ and $\mathcal{YD}(H)$ are isomorphic as braided T -categories over G .*

Remark 3.11 From $\mathcal{YD}(H)$ (respectively $DT(H)$) we can obtain, by pull-back along the group morphism $Aut_{Hopf}(H) \rightarrow G, \alpha \mapsto (\alpha, id)$, a braided T -category (respectively a quasitriangular T -coalgebra) over $Aut_{Hopf}(H)$. If π is a group together with a group morphism $\pi \rightarrow Aut_{Hopf}(H)$, by pull-back along it we obtain a braided T -category (respectively a quasitriangular T -coalgebra) over π . Quasitriangular T -coalgebras over such π have been studied by Virelizier in [17].

4 An isomorphism of categories ${}_H \mathcal{YD}^H(\alpha, \beta) \simeq {}_H \mathcal{YD}^H$ in the presence of a pair in involution

The aim of this section is to prove the following result.

Theorem 4.1 *Let $\alpha, \beta \in Aut_{Hopf}(H)$ and assume that there exists (f, g) a pair in involution corresponding to (α, β) . Then the categories ${}_H \mathcal{YD}^H(\alpha, \beta)$ and ${}_H \mathcal{YD}^H$ are isomorphic. A pair of inverse functors (F, G) is given as follows. If $M \in {}_H \mathcal{YD}^H(\alpha, \beta)$, then $F(M) \in {}_H \mathcal{YD}^H$, where $F(M) = M$ as vector space, with structures*

$$\begin{aligned} h \rightarrow m &= f(\beta^{-1}(S(h_1)))\beta^{-1}(h_2) \cdot m, \\ m \mapsto m_{<0>} \otimes m_{<1>} &= m_{(0)} \otimes m_{(1)}g^{-1}. \end{aligned}$$

If $N \in {}_H \mathcal{YD}^H$, then $G(N) \in {}_H \mathcal{YD}^H(\alpha, \beta)$, where $G(N) = N$ as vector space, with structures

$$\begin{aligned} h \rightharpoonup n &= f(h_1)\beta(h_2) \cdot n, \\ n \mapsto n^{(0)} \otimes n^{(1)} &= n_{(0)} \otimes n_{(1)}g. \end{aligned}$$

Both F and G act as identities on morphisms.

Proof. One checks, by direct computation, that F and G are functors, inverse to each other. Alternatively, we can give a very short proof using results from the previous section. By Example 2.7, we have $fk^g \in {}_H\mathcal{YD}^H(\alpha, \beta)$. By Proposition 3.6, we get $(fk^g)^* \in {}_H\mathcal{YD}^H((\alpha, \beta)^{-1})$. Then, one can check that actually $F(M) = (fk^g)^* \otimes M \in {}_H\mathcal{YD}^H$ and $G(N) = (fk^g) \otimes N \in {}_H\mathcal{YD}^H(\alpha, \beta)$. Also, one can see that $(fk^g)^* \otimes fk^g = fk^g \otimes (fk^g)^* = k$ as objects in ${}_H\mathcal{YD}^H$, hence $F \circ G = G \circ F = id$, using the associativity of the tensor product. $\square$

As we have noticed before, for any $\alpha \in Aut_{Hopf}(H)$ we have that $(\varepsilon, 1)$ is a pair in involution corresponding to (α, α) , hence we obtain:

Corollary 4.2 ${}_H\mathcal{YD}^H(\alpha, \alpha) \simeq {}_H\mathcal{YD}^H$.

Also, as a consequence of the theorem, we obtain the following result (a right-left version was given in [14]), which might be useful for the area of applicability of anti-Yetter-Drinfeld modules:

Corollary 4.3 *Assume that there exists a pair in involution (f, g) corresponding to (S^2, id) . Then the category ${}_H\mathcal{YD}^H(S^2, id)$ of anti-Yetter-Drinfeld modules is isomorphic to ${}_H\mathcal{YD}^H$, and any anti-Yetter-Drinfeld module can be written as a tensor product $fk^g \otimes N$, with $N \in {}_H\mathcal{YD}^H$.*

Let again $\alpha, \beta \in Aut_{Hopf}(H)$ such that there exists (f, g) a pair in involution corresponding to (α, β) , and assume that H is finite dimensional. Then we know that ${}_H\mathcal{YD}^H(\alpha, \beta) \simeq {}_{H^* \bowtie H(\alpha, \beta)}\mathcal{M}$, ${}_H\mathcal{YD}^H \simeq {}_{D(H)}\mathcal{M}$, and the isomorphism ${}_H\mathcal{YD}^H(\alpha, \beta) \simeq {}_H\mathcal{YD}^H$ constructed in the theorem is induced by an algebra isomorphism between $H^* \bowtie H(\alpha, \beta)$ and $D(H)$, given by

$$\begin{aligned} D(H) &\rightarrow H^* \bowtie H(\alpha, \beta), & p \otimes h &\mapsto g^{-1} \rightharpoonup p \bowtie f(\beta^{-1}(S(h_1)))\beta^{-1}(h_2), \\ H^* \bowtie H(\alpha, \beta) &\rightarrow D(H), & p \bowtie h &\mapsto g \rightharpoonup p \otimes f(h_1)\beta(h_2). \end{aligned}$$

References

- [1] D. Bulacu, F. Panaite, F. Van Oystaeyen, Generalized diagonal crossed products and smash products for (quasi) Hopf algebras. Applications, in preparation.
- [2] S. Caenepeel, M. De Lombaerde, A categorical approach to Turaev's Hopf group-coalgebras, arXiv:math.QA/0409600.
- [3] S. Caenepeel, G. Militaru, S. Zhu, "Frobenius and separable functors for generalized module categories and nonlinear equations", *Lecture Notes in Mathematics* **1787**, Springer Verlag, Berlin, 2002.
- [4] S. Caenepeel, F. Van Oystaeyen, Y. Zhang, The Brauer group of Yetter-Drinfeld module algebras, *Trans. Amer. Math. Soc.* **349** (1997), 3737–3771.
- [5] A. Connes, H. Moscovici, Hopf algebras, cyclic cohomology and the transverse index theorem, *Comm. Math. Phys.* **198** (1998), 199–264.
- [6] A. Connes, H. Moscovici, Cyclic cohomology and Hopf algebra symmetry, *Lett. Math. Phys.* **52** (2000), 1–28.
- [7] P. M. Hajac, M. Khalkhali, B. Rangipour, Y. Sommerhauser, Stable anti-Yetter-Drinfeld modules, *C. R. Math. Acad. Sci. Paris* **338** (2004), 587–590.

- [8] P. M. Hajac, M. Khalkhali, B. Rangipour, Y. Sommerhauser, Hopf-cyclic homology and cohomology with coefficients, *C. R. Math. Acad. Sci. Paris* **338** (2004), 667–672.
- [9] F. Hausser and F. Nill, Diagonal crossed products by duals of quasi-quantum groups, *Rev. Math. Phys.* **11** (1999), 553–629.
- [10] P. Jara, D. Ştefan, Cyclic homology of Hopf Galois extensions and Hopf algebras, arXiv:math.KT/0307099.
- [11] C. Kassel, "Quantum groups", *Graduate Texts in Mathematics* **155**, Springer Verlag, Berlin, 1995.
- [12] S. Majid, "Foundations of quantum group theory", Cambridge Univ. Press, 1995.
- [13] S. Montgomery, "Hopf algebras and their actions on rings", CBMS Regional Conference Series, Vol. 82, Amer. Math. Soc., Providence, RI, 1993.
- [14] M. D. Staic, A note on anti-Yetter-Drinfeld modules, preprint 2004.
- [15] M. E. Sweedler, "Hopf algebras", Benjamin, New York, 1969.
- [16] V. Turaev, Homotopy field theory in dimension 3 and crossed group-categories, arXiv:math.GT/0005291.
- [17] A. Virelizier, Graded quantum groups, arXiv:math.QA/0312330.
- [18] M. Zunino, Yetter-Drinfeld modules for crossed structures, *J. Pure Appl. Algebra* **193** (2004), 313–343.